

Year 1998

Amyloid A component

Simcha U., Patrizia C., Shlomit E. and Yaacov M.

Journal of Histochemistry and Cytochemistry; 46: 1377-1384, December 1998.

Widespread expression of serum amyloid A in histologically normal human tissues: predominant localization to the epithelium.

c-erbB-2

Wu J., Zhang P., Lyons B. and Wu L.

Journal of Clinical Laboratory Analysis; 12(5): 298-305, December 1998.

Isolation of the intact molecule and ectodomain of C-erbB-2 oncoprotein from SK-BR-3 cells and development of immunoassays on microplate.

DAB

Mignon B., Nikkels A., Piérard G., Losson B.

Dermatology; 196: 438-441, 1998.

The in vitro and in vivo production of a 31.5-kD keratinolytic subtilase from *Microsporum canis* and the clinical status in naturally infected cats.

Year 1999

Cytokeratin (High Mol. Wt.) (Clone AE3)

Valentin H., Azocar O., Horvat B., Williems R., Garrone R., Evlashev A., Toribio M. and Rabourdin-Combe C.

Journal of Virology; 73(3): 2212-2221, March 1999.

Measles virus infection induces terminal differentiation of human thymic epithelial cells.

Year 2000

Cyclin E

Richter J., Wagner U., Kononen J., Fijan A., Bruderer J., Schmid U., Ackermann D., Maurer R., Alund G., Kno"nagel I., Rist M., Wilber K., Anabitarte M., Hering F., Hardmeier T., Scho"nenberger A., Renata F., Ja"ger P., Fehr J., Schraml P., Moch H., Mihatsch M., Gasser T, Kallioniemi O. and Sauter G.

American Journal of Pathology; 157(3): 787-794, September 2000.

High-throughput tissue microarray analysis of cyclin E Gene amplification and overexpression in urinary bladder cancer.

GST-pi

Tanaka S., Kamitani H., Amin M., Watanabe T., Oka H., Fujii K., Nagashima T. and Hori T.

Journal of Neuro-oncology; 46(2): 157-171, January 2000

Preliminary individual adjuvant therapy for Gliomas based on the results of molecular biological analyses for drug-resistance genes.

p63

Gafa R., Maestri I., Matteuzzi M., Santini A., Ferretti S., Cavazzini L. and Lanza G.
Cancer; 89(10): 2025-2037, November 2000.

Sporadic colorectal adenocarcinomas with high-frequency microsatellite instability
Pathobiological features, hMLH1 and hMSH2 expression, and clinical outcome.

Year 2001

Cytokeratin

Cenacchi G., Roncaroli F., Cerasoli S., Ficarra G., merli G. and Giangaspero F.

The American Journal of Surgical Pathology; 25(3): 401-405, March 2001.

Chordoid glioma of the third ventricle: an ultrastructural study of three case with a histogenetic hypothesis.

Ki-67

Beltrami A., Urbanek K., Kajstura J., Yan S., Finato N., Bussani R., Nadal-Ginard B., Silvestri F., Leri A., Beltrami C. and Anversa P.

The New England journal of Medicine; 344(23): 1750-1757, June 2001.

Evidence that human cardiac myocytes divide after myocardial infarction.

p53

Satoh K., Kihira K., Kawata H., Tokumaru K., Kumakura Y., Ishino Y., Kawakami S., Inoue K., Kojima T., Satoh Y., Mutoh H. and Sugano K.

Helicobacter; 6(1): 31-36, March 2001.

p53 expression in the gastric mucosa before and after eradication of Helicobacter pylori.

Year 2002

CD34

Su M., Tokairin T., Nishikawa Y., Yoshioka T., Takahashi O., Watanabe H., Doi Y., Omori Y., Yoshioka T., Sageshima M., Tanaka T. and Enomoto K.

Pathology International; 52(2): 158-63, February 2002.

Primary osteosarcoma of the uterine corpus: case report and review of the literature.

Cytokeratin 17

Yasuhara Y., Sakaida N., Uemura Y., Senzaki H., Shikata N. and Tsubura A.

Pathology International; 52(4): 158-63, April 2002.

Serous microcystic adenoma (glycogen-rich cystadenoma) of the pancreas: Study of 11 cases showing clinicopathological and immunohistochemical correlations.

Cytokeratin (High Mol. Wt.) (Clone AE3)

Vidalain P., Laine D., Zaffran Y., Azocar O., Servet-Delprat C., Wild T., Rabourdin-Combe C. and Valentin H.

Journal of Virology; 76(13): 6415-6424, July 2002.

Interferons mediate terminal differentiation of human cortical thymic epithelial cells.

Ki-67

Ladage P., Petroll W., Jester J., Fisher S., Begmanson J. and Cavanagh H.
CLAO Journal; 28(4): 177-180, October 2002.

Spherical indentations of human and rabbit corneal epithelium following extended contact lens wear

Macrophage, CD68

Groisman G., Amar M., Schafer I.

Applied Immunohistochemistry & Molecular Morphology; 10(3): 205-209, September 2002.

The Histiocytic marker PG-M1 is helpful in differentiating histiocytes and histiocytic tumors from melanomas.

MPO

Akamizu T., Ozaki S., Hiratani H., Uesugi H., Sobajima J., Hataya Y., Kanamoto N., Saijo M., Hattori Y., Moriyama K., Ohmori K. and Nakao K.

Clinical & Experimental Immunology; 127(1): 92-98, January 2002.

Drug-induced neutropenia associated with anti-neutrophil cytoplasmic antibodies (ANCA): possible involvement of complement in granulocyte cytotoxicity.

P63

Lanza G., Gafa R., Maestri I., Santini A., Matteuzi M. and Cavazzini L.

Modern Pathology; 15(7): 741-749, July 2002.

Immunohistochemical pattern of MLH1/MSH2 expression is related to clinical and pathological features in colorectal adenocarcinomas with microsatellite instability.

Rb

Gu L.; Ichiki Y.; Kitajima Y.

Journal of Cutaneous Pathology; 29(8): 473-479, September 2002.

Aberrant expression of p16 and RB protein in eccrine porocarcinoma.

Synaptophysin

O'Brien R., Xu D., Mi R., Tang X., Hopf C. and Worley P.

The Journal of Neuroscience; 22(11): 4487-4498, June 2002.

Synaptically targeted Narp plays an essential role in the aggregation of AMPA receptors at excitatory synapses in cultured spinal neurons.

Year 2003

Androgen Receptor

Chihara Y., Fujimoto K., Takada S., Hirayama A., Cho M., Yoshida K., Ozono S. and Hirao Y.

International Journal of Urology; 10(12): 672, December 2003.

Aggressive angiofibroma in the scrotum expressing androgen and progesterone receptors.

GFAP

Iseda T., Nishio T., Kawaguchi S., Kawasaki T. and Wakisaka S.

Journal of Comparative Neurology; 464; 343-355, 2003

Nitric Oxide and cGMP mediate alpha1D adrenergic receptor-stimulated protein secretion and p42/p44 MAPK activation in rat lacrimal gland.

GST-pi

Takuya M., Katsuhisa K., Akihito F., Kohei T., Naoki T., Naofumi T., Ikuta T., Takehide A., Tsutomu S., Tetsuji T., Takeshi T., Sumio S. and Yoshiro N.

International Journal of Clinical Oncology; 8(5) :326-331, October 2003.

C(H)OP refractory chronic lymphocytic leukemia patients in whom salvage chemotherapy chosen by evaluating multiple chemotherapeutic drug-resistant factors was remarkably effective.

Ki-67

Ladage P., Jester J., Petroll W., Bergmanson J. and Cavanagh H.

Investigative Ophthalmology & Visual Science; 44(3): 1056-1063, March 2003.

Vertical movement of epithelial basal cells toward the corneal surface during use of extended-wear.

Melanoma

Schuerfeld K., Lazzi S., de Santi M., Gozzetti A., Leoncini L. and Pileri S.

Histopathology; 43(5): 491-494, November 2003.

Cytokeratin-positive interstitial cell neoplasm: a case report and classification issues.

T₄

Nishimura N., Yonemoto J., Miyabara Y., Sato M. and Tohyama C.

Endocrinology, 144(5): 2075-2083, May 2003.

Rat Thyroid hyperplasia induced by gestational and lactational exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin.

Year 2004

Bcl-6

Traggiai E., Chicha L., Mazzucchelli L., Bronz L., Piffaretti J., Lanzavecchia A., and Manz M.

Science; 304 (5667): 104-107, April 2004.

Development of a human adaptive immune system in cord blood cell-transplanted mice.

CD44

Gadalla H., Kamel N., Badary F. and Elanany F.

BJU International; 93(1): 151, January 2004.

Expression of CD44 protein in bilharzial and non-bilharzial bladder cancers.

Cytokeratin 7

Idilman R., Erden E., Kuzu I., Sadik E., Zeki K., Kaan K., Gul Y., Yaman T., Sahin Y., Ajlan T., Ulus A. and Selim K.

Transplantation; 78; 1647-1652, 2004

Recipient-derived hepatocytes in sex-mismatched liver allografts after liver transplant: Early versus late transplant biopsy.

Smooth Muscle Actin

Hodges R., Shatos M., Tarko R., Vrouvlianis J., Gu J. and Dartt D.

Investigative Ophthalmology and Visual Science; 46: 2781-2789, 2005.

Nitric Oxide and cGMP mediate alpha1D adrenergic receptor-stimulated protein secretion and p42/p44 MAPK activation in rat lacrimal gland.

Cytokeratin 17

Pei R., Danbara N., Tsujita-Kyutoku M., Yuri T. and Tsubura A.

Medical Electron Microscopy; 37(2): 114-118, June 2004.

Immunohistochemical profiles of Mallory body by a panel of anti-cytokeratin antibodies.

Ki-67

1) Horackova M., Arora R., Chen R., Armour J., Cattini P., Livingston R., Byczko Z.

American Journal of Physiology. Heart and Circulatory Physiology; 287(4):H1599-608, October 2004.

Cell transplantation for the treatment of acute myocardial infarction: the unique capacity for repair by skeletal muscle satellite cells.

2) Bhattacharya A., Tóth K., Mazurchuk R., Sperryak J., Slocum H., Pendyala L., Azrak R., Cao S., Durrani F. and Rustum Y.

Clinical Cancer Research 10: 8005-8017, December 2004.

Lack of microvessels in well-differentiated regions of human head and neck squamous cell carcinoma A253 associated with functional magnetic resonance imaging detectable hypoxia, limited drug delivery, and resistance to Irinotecan therapy.

MART-1

Morioka M., Hamada J., Ohmori Y., Kageshita T. and Ushio Y.

Neurosurgery; 55(4): 983-984, October 2004.

Spinal arachnoid cyst containing nevus cells in a patient with a large congenital melanocytic nevus: case report.

MMP-14

Suarez-Cuervo C., Merrell M., Watson L., Harris K., Rosenthal E., Väänänen H. and Selander K.

Clinical and Experimental Metastasis; 21(6)-525-533, January 2004.

Breast cancer cells with inhibition of p38 have decreased MMP-9 activity and exhibit decreased bone metastasis in mice.

Year 2005

CD3

Jayakumar P., Berger I., Autschbach F., Weinstein M., Funke B., Verdin E., Goldsmith M. and Keppler O.

Journal of Virology; 79(8): 5220-5226, April 2005.

Tissue-resident macrophages are productively infected ex vivo by primary X4 isolates of human immunodeficiency virus type 1.

Ki-67

Robertson D., Li L., Fisher S., Pearce V., Shay J., Wright W., Cavanagh H. and Jester J.

Investigative Ophthalmology & Visual Science; 46(2): 470, February 2005.

Characterization of growth and differentiation in a telomerase-immortalized human corneal epithelial cell line.

Melanoma

Baldi A., Lombardi D., Russo P., Palescandolo E., De Luca A., Santini D., Baldi F., Rosiello L., Dell'Anna M., Mastrofrancesco A., Maresca V., Flori E., Natali P., Picardo M. and Paggi M.

Clinical Cancer Research; 11: 3175-3183, May 2005.

Ferritin contributes to melanoma progression by modulating cell growth and sensitivity to oxidative stress.

Smooth Muscle Actin

Hodges R., Shatos M., Tarko R., Vrouvlianis J., Gu J. and Dartt D.

Investigative Ophthalmology and Visual Science; 46: 2781-2789, 2005.

Nitric Oxide and cGMP mediate alpha1D adrenergic receptor-stimulated protein secretion and p42/p44 MAPK activation in rat lacrimal gland.

Thrombospondin

Karavasilis V., Mitsi-Malamou V., Briasoulis E., Tsanou E., Kitsou E., Kalofonos H., Fountzilias G., Fotsis T. and Pavlidis N.

BMC Cancer; 5: 25, 2005

Angiogenesis in cancer of unknown primary: Clinicopathology study of CD34, VEGF and TSP-1.

VEGFR-1, VEGFR-2

Kyzas P A., Stefanou D., Baitstaou A. and Agnantis N J.

Modern Pathology; 18: 485-494, 2005

Potential autocrine function of vascular endothelial growth factor in head and neck cancer via vascular endothelial growth factor receptor-2.